# visocolor<sup>®</sup>ECO

**Phosphate** 

en

Test kit for performing colorimetric tests on phosphate ions in surface water and sewage

### Method:

Ammonium molybdate forms with phosphate ions phosphomolybdic acid, which is reduced to phosphomolybdenum blue.

Measurement range: 0.2–5 mg/L PO<sub>4</sub>-P

# Contents of test kit (\*refill pack):

sufficient for 80 tests

- 25 mL PO<sub>4</sub>-1\*
- 25 mL PO<sub>4</sub>-2\*
  - 2 screw-plug measuring glasses
  - 1 slide comparator 1 colour chart
  - 1 plastic syringe 5 mL
  - 1 instructions for use\*

#### Hazard warning:

Information regarding safety can be found on the box' label and in the safety data sheet. You can download the SDS from *www.mn-net.com/SDS*.

#### Instructions for use:

also refer to the pictogram on the back of the color chart

- Pour a 5 mL water sample into each of the measuring glasses using the plastic syringe.
- Place a measuring glass on position A in the comparator.

## Only add the reagent to measuring glass B.

- 2. Add 6 drops of PO<sub>4</sub>-1, seal the glass and mix.
- 3. Add 6 drops of PO<sub>4</sub>-2, seal the glass and mix.
- 4. Open the glass after 10 min and place it on position B in the comparator.
- Slide the comparator until the colors match in the inspection hole on top. Check the measurement reading in the recess on the comparator reed. Mid-values can be estimated.

6. After use, rinse out both measuring glasses thoroughly and seal them. The reagents can be used for the **photometric evaluation** with photometer PF-12/PF-12<sup>Plus</sup>.

This technique can be used also for analyzing sea water.

#### Disposing of the samples:

Information regarding disposal can be found in the safety data sheet. You can download the SDS from *www.mn-net.com/SDS*.

#### Interferences:

Larger amounts of oxidizing reagents inhibit formation of the blue color complex and have to be destroyed.  $H_2S$  interferes in concentrations above 2 mg/L, but can be expelled after acidification of the water sample. Heavy metals in excess of 10 mg/L can slightly decrease the intensity of the color (vanadium causes an increase in color). Silica interferes in excess of 10 mg/L Si.

#### **Conversion table:**

mg/L PO <sub>4</sub> -P	mg/L PO <sub>4</sub> <sup>3-</sup>	mg/L P <sub>2</sub> O <sub>5</sub>
0.2	0.6	0.5
0.3	0.9	0.7
0.5	1.5	1.1
0.7	2.1	1.6
1	3	2
2	6	5
3	9	7
5	15	12

Storage:

Store the test kit in a cool (< 25 °C) and dry place.